

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/329577446>

Gene Expression Profiles of Odorant Binding Proteins in *Glossina brevipalpis* (Diptera: Glossinidae)

Article in *Journal of Entomological Science* · October 2018

DOI: 10.18474/JES17-103.1

CITATIONS

2

READS

138

5 authors, including:



Mary Murithi

Jomo Kenyatta University of Agriculture and Technology

7 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)



Rosaline Wanjiru Macharia

University of Nairobi

32 PUBLICATIONS 399 CITATIONS

[SEE PROFILE](#)



Esther Magiri

Jomo Kenyatta University of Agriculture and Technology

33 PUBLICATIONS 246 CITATIONS

[SEE PROFILE](#)



Steven Nyanjom

Jomo Kenyatta University of Agriculture and Technology

52 PUBLICATIONS 504 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Ecology of arbovirus mosquito vectors [View project](#)



Impact of Genetically modified organisms [View project](#)

Gene Expression Profiles of Odorant Binding Proteins in *Glossina brevipalpis* (Diptera: Glossinidae)

Author(s): Mary Murithi, Rosaline Macharia, Esther Magiri, Steven Nyanjom, and Daniel Masiga

Source: Journal of Entomological Science, 53(4):467-477.

Published By: Georgia Entomological Society

<https://doi.org/10.18474/JES17-103.1>

URL: <http://www.bioone.org/doi/full/10.18474/JES17-103.1>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Gene Expression Profiles of Odorant Binding Proteins in *Glossina brevipalpis* (Diptera: Glossinidae)¹

Mary Murithi², Rosaline Macharia³, Esther Magiri⁴, Steven Nyanjom⁵, and Daniel Masiga⁶

School of Pharmacy, Pharmacy Department, Kabarak University, Private Bag 20157, Kabarak, Kenya

J. Entomol. Sci. 53(4): 467–477 (October 2018)

Abstract *Glossina* spp. (Diptera: Glossinidae) vector trypanosomes, the causal pathogens of trypanosomiasis in humans and animals. *Glossina* spp. chemosensory mechanism is relevant to understanding the behavioral responses of these insects to environmental cues. Odorant binding proteins (OBPs) are involved in either olfactory or nonolfactory cue sensation depending on their spatial and temporal expression. In most insects, the OBPs are ubiquitously expressed, although the levels are higher in the antennae. This study was aimed at determining the differential gene expression patterns of OBP genes in the developmental stages (larval, pupal, teneral) and differentially starved adults of *Glossina brevipalpis* (Newstead) collected from Shimba Hills National Park, Kenya. Five OBP genes, *GbrOBP2*, *GbrOBP6*, *GbrOBP7*, *GbrOBP8*, and *GbrOBP13*, were selected for quantitative PCR expression analysis as representatives of the three OBP classes in *Glossina* spp. (classic, minus-C, classic-dimers). Antennal samples from the newly emerged insects (teneral) served as the controls. *GbrOBP2* was highly expressed in both larval and pupal stages, whereas *GbrOBP7* was expressed in the pupal stage. *GbrOBP8*, *GbrOBP13*, and *GbrOBP6* showed high expression in adult antennae. Expression of OBPs in the immature stages suggested OBP involvement in nonolfactory chemical sensation, whereas that in adult antennae was attributed to olfaction.

Key Words *Glossina brevipalpis*, larvae, pupae, teneral, odorant binding proteins, qPCR

Glossina spp. (Diptera: Glossinidae) are the only known biological vectors of African trypanosomes, the pathogens responsible for human African trypanosomiasis, or sleeping sickness, and animal African trypanosomiasis, or nagana, in livestock (Bouyer et al. 2006). The disease puts approximately 70 million people and 60 million cattle within 37 countries of sub-Saharan Africa at risk of infection (Cecchi and Mattioli 2009, WHO 2013). Additionally, nagana transmitted by *Glossina* spp. lowers agricultural and livestock production, resulting in annual losses estimated at US\$5 billion (Samdi et al. 2011). All *Glossina* spp. are

¹Received 04 September 2017; accepted for publication 11 December 2017.

²Corresponding author (email: mmury@kabarak.ac.ke).

³Centre for Biotechnology and Bioinformatics, University of Nairobi, PO Box 30197-00100, Nairobi, Kenya.

⁴Co-operative University of Kenya, PO Box 24814-00502, Karen, Nairobi, Kenya.

⁵Department of Biochemistry, Jomo Kenyatta University of Agriculture and Technology, PO Box 62000-00200, Nairobi, Kenya.

⁶Molecular Biology and Bioinformatics Unit, International Centre of Insect Physiology and Ecology, ICIPE, PO Box 30772-00100, Nairobi, Kenya.

considered potential vectors of trypanosomes. Control strategies targeting the vector such as sterile insect technique, use of odor-baited traps, and insecticide-treated targets are suitable to reduce the transmission of trypanosomes (Omolo et al. 2009, Abda-Alla et al. 2013). However, the molecular mechanism by which different *Glossina* spp. respond to various hosts is not clearly understood. For instance, Gikonyo et al. (2002) reported a preferential feeding of *Glossina pallidipes* (Austen) toward water buffalo and oxen as opposed to waterbuck.

The study reported herein focused on profiling the expression of putative odorant binding protein (OBP) genes in *Glossina brevipalpis* (Newstead), a species of *Glossina* associated with forested habitats that is known to transmit animal trypanosomiasis (Krafsur 2009). The species occurs mostly in Central and West Africa but is also widely found in Southern and Eastern Africa, with discontinuous occurrence in Mozambique, northern Tanzania, and southeastern Kenya (Krafsur 2009, Cecchi et al. 2008).

Olfactory organs in *Drosophila melanogaster* (Meigen) are the antennae and the maxillary palps, which house about 1,200 and 120 olfactory receptor neurons (ORNs), respectively. The ORNs have sensilla that are divided morphologically into basiconic, trichoid, and coeloconic, and they function in the detection of fruit odors, pheromones, organic acids, and amines, respectively. Once stimulated, the ORNs generate action potential relative to the quality and intensity of the stimulus that activates the higher centers in the brain. The dendrites of up to four ORNs are found in each sensillum (Hallem et al. 2004, Hallem and Carlson 2006). OBPs occur inside the sensilla with their expression varying across the different sensilla classes and cell types. The occurrence of OBPs within the sensilla is abundant and diverse. More so, their expression is sensillum specific. The basiconic, coeloconic, and trichoid sensilla in *Drosophila* were found to express at least one abundant OBP, signifying their relevance in sensilla information coding (Larter et al. 2016).

OBPs are small globular proteins of 120–150 amino acids with molecular weights of 14–16 kDa and contain a signal peptide (Zhou et al. 2008). They are water soluble and extracellular proteins that are secreted by nonneuronal support cells into the sensilla lymph (Kim and Smith 2001). In insects, OBPs are mainly classified into four subfamilies defined by the occurrence of conserved cysteine residues, which influence the tertiary folding (Hekmat-Scafe et al. 2002). These include classic (conserved six cysteine signature), plus-C (more than six cysteines plus a conserved proline), minus-C (lacking the second and the fifth cysteine residues), and classic-dimers (two classical OBPs) (Fan et al. 2011). Zhou et al. (2008) reported that most OBPs belong to the classic subfamily. No plus-C OBPs were identified in *Glossina* spp. (Macharia et al. 2016). Insect OBPs consist of residues that are very divergent within and across species (Pelosi et al. 2017) but with a conserved tertiary structure suggesting a conserved function.

During olfaction, the odorants from the environmental cues (i.e., sex, alarm, or aggregation pheromones) enter into the sensilla through the cuticle pores. While inside, they bind OBPs that transport them to the receptors on the ORNs. Action potential is generated and the signal is carried to the antennal lobe, the primary olfactory center in the brain. The signal is then transmitted to higher brain centers, which include the mushroom bodies and the lateral protocerebrum (Gadenne et al. 2016). This causes them to elicit olfactory-mediated behaviors, (i.e., identification of hosts and mates, evading predators, and location of suitable resting sites) (Vieira

and Rozas 2011, Masiga et al. 2014). The OBPs may also be involved in binding lipophilic odorant molecules or in deactivation and clearing of odors after stimulation (Gong et al. 2009).

Genome annotations have revealed that OBPs vary in number among insects. For instance, reports on other dipterans indicate that there are 51 OBPs in *D. melanogaster* (Hekmat-Scafe et al. 2002), 69 in *Anopheles gambiae* (Say), 111 in *Aedes aegypti* (L.) and 109 in *Culex quinquefasciatus* (Say) (Manoharan et al. 2013), 44 in *Bombyx mori* (L.) (Gong et al. 2009), and 46 in *Tribolium castaneum* (Herbst) and 21 in *Apis mellifera* (L.) (Forêt and Maleszka 2006). In *Glossina* spp., the number of OBPs are as follows: 30 in *G. morsitans morsitans* (Westwood), *G. fuscipes* (Newstead) as well as in *G. pallidipes*, and 29 in *G. austeni* (Newstead) and 28 in *G. brevipalpis* (Macharia et al. 2016). The low number of OBPs in *Glossina* compared with other insects could be attributed to the fact that they exclusively feed on blood and with some degree of host preferences, which may have resulted to streamline their chemoreception (International Glossina Genome Initiative 2014).

The expression of OBPs in nonolfactory organs of insects, such as legs, wings, fat body, ovaries, and pupae, has been attributed to possible involvement in nonolfactory functions (Pelosi et al. 2017). For instance, significant expression of OBP45 in *Ae. aegypti* female ovaries at 48 h post-blood meal (pbm) suggested its probable role in egg maturation during reproduction (Costa-da-Silva et al. 2013).

This study presents, for the first time to best of our knowledge, the expression analysis of five OBP genes at different developmental stages and differentially starved adults of *G. brevipalpis*. The findings provide useful insights into the *Glossina* spp. biology of chemoreception that may be exploited in designing specific vector control strategies.

Materials and Methods

Sample collection. Adult *G. brevipalpis* were collected from Shimba Hills National Park, located in Kwale County, Kenya (E 04°14', S 039°23', elevation 445 m above sea level) in December 2014 by using biconical traps baited with acetone, 1-octen-3-ol, and 4-methylphenol (Challier and Laveissiere 1973). The *G. brevipalpis* specimens were identified morphologically from other *Glossina* spp. The insects were maintained exclusively on rabbit blood in accordance with standard procedures for animal care in the insectary at the International Center of Insect Physiology and Ecology (ICIPE) from where the larva and pupa were obtained. Antennae from female teneral and adults starved at periods of 2, 24, 48, 72, and 96 h were dissected on ice, and the samples were preserved in liquid nitrogen until used for RNA extraction.

Total RNA extraction. Total RNA from the larval, pupal, and antennal samples was extracted using TRIzol (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The presence of any contaminating genomic DNA was digested with RNase free-DNase1 (Fermentas, Life Sciences, Waltham, MA, USA). The purity of the RNA obtained was assessed using a spectrophotometer (Nanodrop 2000, Thermo Scientific, Waltham, MA, USA) and integrity was

Table 1. Primers for the five putative *G. brevipalpis* OBP genes that were quantified using qRT-PCR.

| Primer Name | Nucleotide Sequence (5' to 3') | |
|------------------|--------------------------------|----------------------|
| | Forward | Reverse |
| GAPDH* | AGATGGAAGCATGTTGCGTA | TATTGGGCCTCCTGCAGATA |
| <i>Gbr</i> OBP2 | ATGGCAGTATCCCGAAGATG | CGCACTTTGCAATCTTTTCA |
| <i>Gbr</i> OBP6 | CTGGCAAGGGAATGCACTAT | CTTTTTGCATGTTCCACAGC |
| <i>Gbr</i> OBP7 | TGGACCAGTGTTTAGCACCA | TTATCAAAGCCCTGCGTCTC |
| <i>Gbr</i> OBP8 | AAGACCGGTACCACAGAGGA | CAATGTGTTGAACGGTTTCC |
| <i>Gbr</i> OBP13 | ACTTAACTCATCGCCCAACG | CGTGAACTGCTGCAAACATT |

* Positive control gene.

determined by running the RNA samples on 1.2% formaldehyde agarose gel electrophoresis.

Synthesis of cDNA. Total RNA from each sample was reverse transcribed in a 20- μ l reaction to a cDNA concentration of 20 ng/ μ l by using the RevertAid First Strand cDNA synthesis kit (Fermentas, Thermo Scientific) following the manufacturer's instructions. The negative control consisted of all the components minus the enzyme reverse transcriptase (RT).

Conventional PCR. Sequences of the putative *G. brevipalpis* OBP genes were downloaded from VectorBase (Giraldo-Calderon et al. 2014) and used in designing primers with the Primer3Plus tool (Untergasser et al. 2012). Amplification of the 28 putative *G. brevipalpis* OBPs was conducted using conventional PCR in 10- μ l reactions. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) from *G. brevipalpis* was used as the internal reference gene. The thermocycling conditions comprised of initial denaturation at 98°C for 30 s, 40 cycles of; denaturation at 98°C for 15 s, annealing at 55°C for 30 s, and extension at 72°C for 30 s, followed by final extension at 72°C for 5 min. The reactions were conducted using a Proflex thermocycler (Applied Biosystems; Thermo Fischer Scientific, Waltham, UK).

Analysis of amplified putative OBPs by using quantitative real-time PCR (qRT-PCR). Quantitative real-time PCR analysis of the PCR-positive putative OBP genes was conducted using a Stratagene Mx3000P qPCR system (Agilent Technologies UK Ltd, Cheshire, UK) in reaction volumes of 10 μ l. The *G. brevipalpis* GAPDH gene was used as the endogenous reference gene. Each reaction contained final concentrations of the following: 1 \times Maxima SYBR Green/ROX master mix (Fermentas, Thermo Scientific, UK), 2 ng/ μ l of cDNA template, 0.3 mM of forward and reverse primers of the putative OBP genes (Table 1), and nuclease-free water. The thermocycling conditions were programmed as follows: one cycle at 95°C for 10 mins, followed by 40 cycles of denaturation at 95°C for 15 s, annealing at 60°C for 30 s, and extension at 72°C for 30 s. A dissociation curve analysis generated at the end of the PCR was used to test for the specificity of the amplification to target genes. All the experimental tests were carried out in triplicate

and a no template control included as a negative control for each set up. The expression changes of each putative OBP gene across the developmental stages and in the differentially starved adults relative to GAPDH were calculated by the $2^{-\Delta\Delta CT}$ (Livak) method (Livak and Schmittgen 2001). Tests of significance of the OBP expression were performed using a single factor analysis of variance (ANOVA), followed by Tukey's test with a threshold of $P < 0.05$.

Results

Expression patterns of five putative OBP genes in the developmental stages of *G. brevipalpis*. Five OBPs were selected for expression analysis based on their linear quantification amplification curve and as representatives of the three OBP classes in *Glossina* spp., which include minus-C OBPs (*GbrOBP2*), dimer OBPs (*GbrOBP7*), and classic OBPs (*GbrOBP6*, *GbrOBP8* and *GbrOBP13*). Of the five genes tested, the gene expression levels of *GbrOBP2* relative to its expression in the control (teneral) were highest in the larval stage, followed by *GbrOBP7* in the pupal stage. The other three, *GbrOBP6*, *GbrOBP8*, and *GbrOBP13*, had minimal expression levels less than that of the control (Fig. 1). The ANOVA P values of *GbrOBP2*, *GbrOBP6*, *GbrOBP7*, *GbrOBP8*, and *GbrOBP13* were $1.59E-06$, $8.32E-07$, $1.13E-07$, $3.59E-07$, and $5.37E-07$ respectively.

Expression patterns of five putative OBP genes in teneral and adult female *G. brevipalpis* starved at different time periods. Out of the five genes, *GbrOBP8* and *GbrOBP13* exhibited the highest expression levels relative to the control (teneral). Expression of *GbrOBP13* peaked at the 72 and 96 h, whereas that of *GbrOBP8* peaked at 24-h postfeeding. Next in abundance were *GbrOBP2* and *GbrOBP6*, with peaked expressions between 24 and 72 h for the former and 24-h postfeeding for the latter. Generally, *GbrOBP7* had the least expression, with lower levels than that of the teneral. Furthermore, the levels remained low for the remainder of the period up to the 96 h. The patterns observed for *GbrOBP6*, *GbrOBP8*, and *GbrOBP13* were similar. Soon after the insects were fed (2-h postfeeding), the expression of the three genes decreased relative to teneral levels, and later in the 24-h period, their expressions increased, followed by a decrease at 48 h and, consequently, a gradual increase from 72 to 96 h (Fig. 2). The ANOVA P values of *GbrOBP2*, *GbrOBP6*, *GbrOBP7*, *GbrOBP8*, and *GbrOBP13* were $7.43E-08$, $2.42E-09$, $1.83E-11$, $3.08E-10$, and $2.15E-12$, respectively.

Discussion

An insect's age influences odor coding. Therefore, there is a variation in their response toward odors emanating from various sources. In its early life, an insect is more attracted to food odors, and then later, it may respond to oviposition-site odors and other odors that are important for its biology. For instance, immediately after emergence, a young female mosquito will seek sugars and later will start searching for vertebrates for blood sucking (Gadenne et al. 2016). Therefore, chemical sensing influences responses in both adults and juvenile insects. Despite this, expression analysis of OBP genes in most insects have only focused on the adults. For instance, information on adult OBPs is available on the lepidopterans




Fig. 1. Expression profiles of five putative OBPs in the larval, pupal ($n = 1$), and teneral (control, $n = 10$) stages of *G. brevipalpis*. All data represent mean values and error bars indicate standard error of the means at 95% confidence. The letters on the bars indicate statistically different expression levels as calculated by the Tukey's test.

Helicoverpa armigera (Hübner) and *H. assulta* (Guenée) (Zhang et al. 2015) and dipterans *Bactrocera dorsalis* (Hendel) (Zheng et al. 2013), *Cx. pipiens* (L.), and *Cx. quinquefasciatus* (Peiletier and Leal 2009)). However, it is also evident that OBP expression in the insect juvenile stage is important, as reported in some studies. Studies on *Chrysomya megacephala* (F.) identified one OBP, *Cmeg33593_c0*, to be highly expressed in the larval stage and was suggested to be involved in larval aging, nutrient accumulation, and regulation of the larval feeding behavior (Wang et al. 2015). In another study with the wheat midge *Sitodiplosis mosellana* (Gehin), the high expression of *SmosOBP1* and *SmosOBP10* in larvae suggested their involvement in larval-specific chemical sensing (Gong et al. 2014). Also, high




Fig. 2. Expression profiles of five putative OBPs in the antenna of *G. brevipalpis* tenerals (control, $n = 10$) and differentially starved adults ($n = 10$). All data represent mean values and error bars indicate standard error of the means at 95% confidence. The letters on the bars indicate statistically different expression levels as calculated by the Tukey's test.

expression of *SexiOBP13* in the larvae of *Spodoptera exigua* (Hübner) suggested its involvement in larval olfaction (Jin et al. 2015).

Unlike most other insects whose larvae are free living, those of *Glossina* spp. are characterized by adenotrophic viviparity (larvae develop *in utero*) whereby the larva obtains its food from the mother's milk glands. On the other hand, the pupal stage, which develops minutes after the larvae have been larviposited, is mainly dormant and is nourished by the fat body (International Glossina Genome Initiative 2014).

The high expression of *GbrOBP2* and *GbrOBP7* in the larval and pupal stages, respectively, in *G. brevipalpis* was similar to that of their orthologs in *G. morsitans morsitans*. These OBPs may, therefore, be suggested to be involved in nonolfactory chemical sensation in the juvenile stages of *Glossina* spp., owing to the fact that the larval and pupal stages have no developed antennae (Liu et al. 2010, 2012). Elsewhere, high expression of *OBP1* in *Anopheles stephensi* (Liston) (*AsteOBP1*) at the pupal stage was attributed to the physiological development of the chemosensory tissue (Sengul and Tu 2010). In yet another study with *A. mellifera* (L.), two OBPs (*OBP10* and *OBP13*) were highly expressed throughout the pupal stages and suggested to be involved in its development while preparing to emerge (Forêt and Maleszka 2006).

Olfaction in adult insects occurs via the well-structured antennae (Jin et al. 2015). Therefore, the expression of OBPs in the antennae of *G. brevipalpis* adults as well as in other insects suggests their involvement in olfaction (Liu et al. 2010, Zheng et al. 2013). Insect feeding state influences their attraction to food whereby, when starved, they become easily attracted to food odors and, after feeding, the insects carry on with other activities relevant for their survival. Therefore, odor-guided behavior, which is driven by peripheral odor-detection proteins, such as OBPs, is driven by the nutritional state of insects. For instance, starved *D. melanogaster* were found to be easily attracted to food odor sources (Gadenne et al. 2016). The nutritional status in *G. morsitans morsitans* was found to affect the expression of antennal OBPs (Otter et al. 2017).

The high expression of *GbrOBP8* in the adults agreed with the findings of Liu et al. (2010) who observed the orthologue gene in *G. morsitans morsitans* (*GmmOBP8*) to be highly expressed in adults and suggested its involvement in olfaction. Differences were also noted. For instance, although *GbrOBP13* was relatively highly expressed in *G. brevipalpis*, this did not correspond with the findings in *G. morsitans morsitans* where the orthologous gene *GmmOBP13* was not among the 10 expressed genes in the antenna. This difference may be attributed to the difference in species host preference. Functional studies of the orthologue of *GbrOBP13* in *D. melanogaster* (*DmelOBP56h*) showed its role in clearing bitter tastes and improving feeding (Swarup et al. 2014). The *GbrOBP6* elevated expression at 24-h pbm and decreased expression at 48-h pbm resembled that of *GmmOBP6* in *G. morsitans morsitans* (Liu et al. 2010). In functional studies of the *GbrOBP6* orthologous gene in *Drosophila*, *DmelOBP28a* showed its involvement in shuttling bitter tastes to odorant receptors, hence discouraging their feeding on bitter foods (Swarup et al. 2014). Therefore, *DmelOBP56h* and *DmelOBP28a* in *D. melanogaster* have been suggested to have opposing functions. Minimal expression of *GbrOBP7* in the antennae of the differentially starved insects agreed with the findings of Liu et al. (2010) whereby the orthologous genes *GmmOBP7* subsequently had low expression levels in the adult antennae of *G. morsitans morsitans*. The variation in the expression of the OBP genes may be due to the different physiological needs of *Glossina* spp. However, it was difficult to pin point the specific roles, as the insects used were field collected and posed a challenge of different variations. However, the findings highlight the pattern of the expression of OBP genes that might be used as a basis in functional characterization. Also, although the insect olfactory system is better studied in adults compared with early developmental stages, this study extends the growing

recognition of the involvement of OBPs in nonolfactory chemical sensing, as depicted in the larval and pupal stages of *Glossina* spp.

Acknowledgments

We are grateful to Lilian Wambua and Vincent Owino for their useful inputs in qPCR data analysis and interpretation. We also appreciate Jeremy Herren, Merid Getahun, and Jackson Muema for valuable edits to the manuscript. Regards to James Kabii for helping with the planning of field logistics. Much regards to the World Federation of Scientists (WFS) for financial support to MM. This work was supported by funding from Jomo Kenyatta University of Agriculture and Technology (grant number JKU/2/4/RP/139).

References Cited

- Abda-Alla, A.M.M., M. Bergoin, A.G. Parker, N.K. Maniana, J.M. Viak, K. Bourtzis and S. Askoy. 2013.** Improving sterile insect technique (SIT) for tsetse flies through research on their symbionts and pathogens. *J. Invertebr. Pathol.* 29: 997–1003.
- Bouyer, J., L. Guerrini, M. Desquesnes, S. de la Rocque and D. Cuisance. 2006.** Mapping African animal trypanosomiasis risk from the sky. *Vet. Res.* 37: 633–45.
- Cecchi, G. and R.C. Mattioli. 2009.** Global geospatial datasets for African trypanosomiasis management: A review, Pp. 1–39. *In* Cecchi G. and R.C. Mattioli RC (ed). Geospatial datasets and analyses for an environmental approach to African trypanosomiasis. Food and Agriculture Organization of the United Nations, Rome.
- Cecchi, G., R.C. Mattioli, J. Slingenbergh and S. De La Rocque. 2008.** Land cover and tsetse fly distributions in sub-Saharan Africa. *Med. Vet. Entomol.* 22: 364–373.
- Challier, A. and C. Laveissiere. 1973.** Un nouveau piege pour la capture des glossines (*Glossina*: Diptera, Muscidae): description et essais sur le terrain. *Ent. Méd. Parasitol.* 9: 251–262.
- Costa-da-Silva, A.L., B.B. Kojin, O. Marinotti, A.A. James and M.L. Capurro. 2013.** Expression and accumulation of the two-domain odorant-binding protein AegOBP45 in the ovaries of blood-fed *Aedes aegypti*. *Parasit. Vectors.* 6: 364.
- Fan, J., F. Francis, Y. Liu, J.L. Chen and D.F. Cheng. 2011.** An overview of odorant-binding protein functions in insect peripheral olfactory reception. *Genet. Mol. Res.* 10: 3056–69.
- Forêt, S. and R. Maleszka. 2006.** Function and evolution of a gene family encoding odorant binding-like proteins in a social insect, the honey bee (*Apis mellifera*). *Genome Res.* 16: 1404–1413.
- Gadanne, C., R.B. Barrozo and S. Anton. 2016.** Plasticity in insect olfaction: To smell or not to smell? *Annu. Rev. Entomol.* 61: 317–33.
- Giraldo-Calderon, G.I., S.J. Emrich, R.M. MacCallum, G. Maslen, E. Dialynas, P. Topalis, H. Nicholas, S. Gesing, G. Madey, F.H. Collins and D. Lawson. 2014.** VectorBase: An updated bioinformatics resource for invertebrate vectors and other organisms related with human diseases. *Nucleic Acids Res.* 43: D707–D713.
- Gong, D.P., H.-J. Zhang, P. Zhao, Q.-Y. Xia and Z.-H. Xiang. 2009.** The odorant binding protein gene family from the genome of silkworm, *Bombyx mori*. *BMC Genomics.* 10: 332.
- Gong, Z., J. Miao, Y. Duan, Y. Jiang, T. Li and Y. Wu. 2014.** Identification and expression profile analysis of putative odorant-binding proteins in *Sitodiplosis mosellana* (Gehin) (Diptera: Cecidomyiidae). *Biochem. Biophys. Res. Commun.* 444: 164–170.
- Hallem, E.A. and J.R. Carlson. 2006.** Coding of odors by a receptor repertoire. *Cell.* 125: 143–160.
- Hallem, E.A., M.G. Ho and J.R. Carlson. 2004.** The molecular basis of odor coding in the *Drosophila* antenna. *Cell.* 117: 965–979.
- Hekmat-Safe, D.S., C.R. Scape, A.J. McKinney and A. Mark. 2002.** Genome-wide analysis of the odorant-binding protein gene family in *Drosophila melanogaster*. *Genome Res.* 12: 1357–1369.

- International Glossina Genome Initiative. 2014.** Genome sequence of the tsetse fly (*Glossina morsitans*): Vector of African trypanosomiasis. *Science*. 344: 207–213.
- Jin, R., N. Liu, Y. Liu and S. Dong. 2015.** A larval specific OBP able to bind the major female sex pheromone component in *Spodoptera exigua* (Hübner). *J. Integr. Agric.* 14: 1356–1366.
- Kim, M.S. and D.P. Smith. 2001.** The invertebrate odorant-binding protein LUSH is required for normal olfactory behavior in *Drosophila*. *Chem. Senses*. 26: 195–199.
- Krafsur, E.S. 2009.** Tsetse flies: Genetics, evolution, and role as vectors. *Infect. Genet. Evol.* 9: 124–141.
- Larter, N.K., J.S. Sun, J.R. Carlson, N. Haven and U. States. 2016.** Organization and function of *Drosophila* odorant binding proteins. *eLIFE Res. Artic.* 5: 1–22.
- Liu, R., X. He, S. Lehane, M. Lehane, M. Berriman, L.M. Field and J. Zhou. 2012.** Expression of chemosensory proteins in the tsetse fly *Glossina morsitans morsitans* is related to female host-seeking behaviour 21: 41–48.
- Liu, R., S. Lehane, X. He, M. Lehane and C.H.M. Berriman. 2010.** Characterisations of odorant-binding proteins in the tsetse fly *Glossina morsitans morsitans*. *Cell. Mol. Life Sci.* 67: 919–929.
- Livak, K.J. and T.D. Schmittgen. 2001.** Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻(Delta Delta C(T)) method. *Methods* 25: 402–408.
- Macharia, R., P. Mireji, E. Murungi, G. Murilla, A. Christoffels, S. Aksoy and D. Masiga. 2016.** Genome-wide comparative analysis of chemosensory gene families in five tsetse fly species. *PLoS Negl. Trop. Dis.* 10: 1–30.
- Manoharan, M., M.N.F. Chong, A. Vaïtinadapoulé, E. Frumence, R. Sowdhamini and B. Offman. 2013.** Comparative genomics of odorant binding proteins in *Anopheles gambiae*, *Aedes aegypti*, and *Culex quinquefasciatus*. *Genome Biol. Evol.* 5: 163–80.
- Masiga, D., G. Obiero, R. Macharia, P. Mireji and A. Christoffels. 2014.** Chemosensory receptors in tsetse flies provide link between chemical and behavioural ecology. *Trends Parasitol.* 30: 1–3.
- Omolo, M.O., A. Hassanali, S. Mpiana, J. Esterhuizen, J. Lindh, M.J. Lehane, P. Solano, J.B. Rayaisse, G.A. Vale, S.J. Torr and I. Tirados. 2009.** Prospects for developing odour baits to control *Glossina fuscipes* spp., the major vector of human African trypanosomiasis. *PLoS Negl. Trop. Dis.* 3: e435.
- Otter, C.J.D., T. Tchicaya and A.M. Schutte. 2017.** Effects of age, sex and hunger on the antennal olfactory sensitivity of tsetse flies. *Physiol. Entomol.* 16: 173–182.
- Peiletier, J. and W.S. Leal. 2009.** Genome analysis and expression patterns of odorant-binding proteins from the Southern House mosquito *Culex pipiens quinquefasciatus*. *PLoS One* 4: e6237.
- Pelosi, P., I. Iovinella, J. Zhu, G. Wang and F.R. Dani. 2017.** Beyond chemoreception: Diverse tasks of soluble olfactory proteins in insects. *Biol. Rev. Camb. Philos. Soc.* 93: 184–200.
- Samdi, S.M., A.O. Fajinmi, J.O. Kalejaye, B. Wayo, M.K. Haruna, J.E. Yarnap, W.P. Mshelia, A.O. Usman, S.M. Hamra, A. Jijitar, R. Ogunwole, R.P. Ovbagbedia and R. Bizi. 2011.** Prevalence of trypanosomiasis in cattle at slaughter in Kaduna central Abattoir. *Asian J. Anim. Sci.* 5: 162–165.
- Sengul, M.S. and Z. Tu. 2010.** Identification and characterization of odorant-binding protein 1 gene from the Asian malaria mosquito, *Anopheles stephensi*. *Insect Mol. Biol.* 19: 49–60.
- Swarup, S., T.V. Morozova, S. Sridhar, M. Nokes and R.R.H. Anholt. 2014.** Modulation of feeding behavior by odorant-binding proteins in *Drosophila melanogaster*. *Chem. Senses* 39: 125–132.
- Untergasser, A., I. Cutcutache, T. Koressaar, J. Ye, B.C. Faircloth, M. Remm and S.G. Rozen. 2012.** Primer3-new capabilities and interfaces. *Nucleic Acids Res.* 40: 1–12.
- Vieira, F.G. and J. Rozas. 2011.** Comparative genomics of the odorant-binding and chemosensory protein gene families across the arthropoda: Origin and evolutionary history of the chemosensory system. *Genome Biol. Evol.* 3: 476–490.

- Wang, X., M. Xiong, C. Lei and F. Zhu. 2015.** The developmental transcriptome of the synanthropic fly *Chrysomya megacephala* and insights into olfactory proteins. *BMC Genomics*. 16: 1–9.
- WHO. 2013.** Control and surveillance of human African trypanosomiasis. World Health Organization technical report series.
- Zhang, J., B. Wang, S. Dong, D. Cao, J. Dong, W.B. Walker, Y. Liu and G. Wang. 2015.** Antennal transcriptome analysis and comparison of chemosensory gene families in two closely related noctuidae moths, *Helicoverpa armigera* and *H. assulta*. *PLoS One*. 10: e0117054.
- Zheng, W., W. Peng, C. Zhu, Q. Zhang, G. Saccone and H. Zhang. 2013.** Identification and expression profile analysis of odorant binding proteins in the oriental fruit fly *Bactrocera dorsalis*. *Int. J. Mol. Sci.* 14: 14936–49.
- Zhou, J.-J., X.-L. He, J.A. Pickett and L.M. Field. 2008.** Identification of odorant-binding proteins of the yellow fever mosquito *Aedes aegypti*: Genome annotation and comparative analyses. *Insect Mol. Biol.* 17: 147–63.